## **SINGAPORE**

# Implementation date and additional rounds 2004, 2006, 2013

#### Package coverage

Health warnings are required to cover 50% of the front and 50% of the back of the package. Overall, 50% of the package space is appropriated to health warnings.

#### **Rotation schedule and history**


Warning labels must appear on an equal number

of packages over the course of a 24-month period. Singapore has produced three rounds of health warning labels. The first set of 6 health warnings was used from 2004 to 2005. A second set of 6 health warnings was released in 2006. A third set of six health warnings is expected to be implemented in 2013.

**Restrictions on misleading information** Unspecified.

### **WARNINGS**










### **WARNINGS**


2006







WARNING: SMOKING CAUSES 92% OF ORAL CANCERS QUIT: 1800-438-2000







WARNING: SMOKING CAUSES GANGRENE QUIT: 1800-438-2000

Tobacco Labelling Resource Centre: Singapore. Waterloo, Ontario: Department of Health, University of Waterloo; 2010 [cited 2011 April 5]; Available from: http://www.tobaccolabels.ca/healthwarningimages/country/singapore.

## **WARNINGS**

2004




WARNING: TOBACCO SMOKE CAN KILL BABIES QUIT: 1800-438-2000



WARNING: SMOKING HARMS YOUR FAMILY QUIT: 1800-438-2000



WARNING: SMOKING CAN CAUSE A SLOW PAINFUL DEATH QUIT: 1800-438-2000



WARNING: SMOKING CAUSES LUNG CANCER QUIT: 1800-438-2000



WARNING: SMOKING CAUSES STROKE QUIT: 1800-438-2000

