

En 2001, une directive du Parlement Européen et du Conseil relative à l'utilisation de mises en garde sanitaires sur les paquets de cigarettes est devenue loi. Les emballages des produits du tabac doivent, au minimum, comporter des mises en garde sous forme de texte ; l'utilisation d'étiquettes de mise en garde sous forme d'illustrations graphiques est facultative. La Commission Européenne (CE) a produit des étiquettes de mise en garde sous forme d'illustrations graphiques et de texte pouvant être utilisées par les États membres.

## Date de mise en œuvre et lots supplémentaires 2004

### Pourcentage du paquet couvert

Les États membres de l'UE doivent impérativement utiliser des mises en garde sous forme de texte conformément aux normes ci-dessous:

#### OBLIGATION MINIME

	sans bordure			avec bordure		
	total	avant	dos	total	avant	dos
Pays unilingues	35%	30%	40%	48%	43%	53%
Pays bilingues	39%	32%	45%	52%	45%	58%
Pays trilingues	43%	35%	50%	56%	48%	63%

## Calendrier de rotation et historique

Une série facultative de 42 mises en garde sous forme d'illustrations graphiques est disponible. La fréquence de rotation recommandée n'est pas précisée.

## Restrictions relatives aux informations trompeuses


L'utilisation de termes descriptifs trompeurs comme « légère » et « douce » est interdite sur les emballages.

## Remarques particulières

Nombre d'États qui ne sont pas membres de l'Union Européenne ont également adopté l'utilisation des images des étiquettes de mises en garde de l'UE sur leurs propres paquets de cigarettes.

# ÉTIQUETTES DE MISE EN GARDE

2004




Tobacco Labelling Resource Centre: European Union. Waterloo, Ontario: Department of Health, University of Waterloo; 2010 [cited 2011 April 6]; Available from: <http://www.tobaccolabels.ca/healthwarningimages/country/europeanunion>.

ÉTIQUETTES DE MISE EN GARDE

2004



Smoking can **damage the sperm** and **decreases fertility**



Smoking can **damage the sperm** and **decreases fertility**



Smoke contains **benzene, nitrosamines, formaldehyde and hydrogen cyanide**

The risk of having a **stroke is approximately 50% higher in smokers than in non-smokers**

Smoking **blocks the arteries** and **causes heart attacks and strokes**



Smoking when pregnant **harms your baby**



Smokers **die younger**



Smoking can cause a **slow and painful death**



Get help to stop smoking:  
**0803 00 00 00**



Smoking causes **fatal lung cancer**




Smoking causes **fatal lung cancer**



Smoking causes **fatal lung cancer**

Stopping smoking **reduces the risk of fatal heart and lung diseases**

Smoking may **reduce the blood flow** and **causes impotence**



Smoking may **reduce the blood flow** and **causes impotence**



Smoking may **reduce the blood flow** and **causes impotence**



Smoking when pregnant **harms your baby**

**You can do it, we can help**

Your **doctor or your pharmacist** can help you stop smoking



Smoking causes **ageing of the skin**



Stopping smoking **reduces the risk of fatal heart and lung diseases**



Smoking causes **ageing of the skin**

**Lifetime smokers lose an average of 14 years of life**

Smokers **die younger**



Smokers **die younger**

**0803 00 00 00**

**Choose freedom, we'll help you**

Get help to stop smoking

**Smoking is a serious nicotine addiction, don't be afraid to ask for help**

Your **doctor or your pharmacist** can help you stop smoking

ÉTIQUETTES DE MISE EN GARDE

2004

